

Sciurid phylogeny and the paraphyly of Holarctic ground squirrels (*Spermophilus*)

Matthew D. Herron, Todd A. Castoe, and Christopher L. Parkinson*

Department of Biology, University of Central Florida, 4000 Central Florida Blvd., Orlando, FL 32816-2368, USA

Received 26 May 2003; revised 11 September 2003

Abstract

The squirrel family, Sciuridae, is one of the largest and most widely dispersed families of mammals. In spite of the wide distribution and conspicuousness of this group, phylogenetic relationships remain poorly understood. We used DNA sequence data from the mitochondrial cytochrome *b* gene of 114 species in 21 genera to infer phylogenetic relationships among sciurids based on maximum parsimony and Bayesian phylogenetic methods. Although we evaluated more complex alternative models of nucleotide substitution to reconstruct Bayesian phylogenies, none provided a better fit to the data than the GTR + G + I model. We used the reconstructed phylogenies to evaluate the current taxonomy of the Sciuridae. At essentially all levels of relationships, we found the phylogeny of squirrels to be in substantial conflict with the current taxonomy. At the highest level, the flying squirrels do not represent a basal divergence, and the current division of Sciuridae into two subfamilies is therefore not phylogenetically informative. At the tribal level, the Neotropical pygmy squirrel, *Sciurillus*, represents a basal divergence and is not closely related to the other members of the tribe Sciurini. At the genus level, the sciurine genus *Sciurus* is paraphyletic with respect to the dwarf squirrels (*Microsciurus*), and the Holarctic ground squirrels (*Spermophilus*) are paraphyletic with respect to antelope squirrels (*Ammospermophilus*), prairie dogs (*Cynomys*), and marmots (*Marmota*). Finally, several species of chipmunks and Holarctic ground squirrels do not appear monophyletic, indicating a need for reevaluation of alpha taxonomy.

© 2003 Elsevier Inc. All rights reserved.

Keywords: Bayesian phylogeny; cytochrome *b*; Marmotini; Sciuridae; *Spermophilus*; Squirrels; Taxonomy

1. Introduction

The squirrel family, Sciuridae, is one of the most diverse and widely distributed families of mammals. With more than 270 species belonging to 50 genera, members of this family occur on all continents except Antarctica and Australia (Hoffmann et al., 1993; McKenna and Bell, 1997). Most species are diurnal and are conspicuous members of the diverse array of ecological communities they inhabit. In spite of the significance of this group, phylogenetic relationships within the family remain poorly understood. Substantial questions remain regarding the monophyly of genera, the relationships among genera, and the validity of the current taxonomy at virtually all levels. In particular, the species content

and monophyly of many recognized genera have not been comprehensively evaluated.

Two subfamilies of squirrels are currently recognized: Pteromyinae, the flying squirrels; and Sciurinae, consisting of nine tribes of arboreal and terrestrial squirrels (McKenna and Bell, 1997). Multiple independent origins have been proposed for the Pteromyinae based on fossil and immunological data (Black, 1963; Engesser, 1979; Hight et al., 1974; James, 1963; Mein, 1970). Studies based on nucleotide and anatomical data, however, have suggested monophyly of the subfamily (Mercer and Roth, 2003; Oshida et al., 1996; Stepan et al., in press; Thorington, 1984). A close relationship between the flying squirrels (whether monophyletic or polyphyletic) and the tribe Sciurini has been inferred based on analyses of fossil and DNA data (Black, 1963, 1972; Mercer and Roth, 2003; Oshida et al., 1996; Stepan et al., in press). Several other studies, though, have supported an early divergence between the

* Corresponding author. Fax: 1-407-823-5769.

E-mail address: cparkins@pegasus.cc.ucf.edu (C.L. Parkinson).

Pteromyinae and the remaining sciurids (Ellis and Maxson, 1980; Engesser, 1979; Hafner, 1984; Hafner et al., 1994; Hight et al., 1974; James, 1963; Forsyth Major, 1893; Mein, 1970; Wade and Gilbert, 1940).

Within the Sciurinae, the tribe Marmotini includes the antelope squirrels (*Ammospermophilus*), prairie dogs (*Cynomys*), marmots (*Marmota*), and Holarctic ground squirrels (*Spermophilus*). Based on fossil data, Bryant (1945) suggested that the chipmunks (*Eutamias*, *Neotamias*, and *Tamias*) were close relatives of the Marmotini, and Moore (1959) considered them a subtribe (Tamiina) of the Marmotini. Later, fossil evidence led Black (1963) to elevate chipmunks to tribal status (Tamiini). Historically, chipmunks have been variously assigned to one genus, sometimes with three subgenera (Bryant, 1945; Ellerman et al., 1940; Levenson et al., 1985; Nadler et al., 1969, 1977); two genera (Ellis and Maxson, 1979; Hall, 1981; Hall and Kelson, 1959; Nadler, 1964; White, 1953); or three genera (Jameson, 1999; Piaggio and Spicer, 2001).

The Marmotini and Tamiini were traditionally thought to be closely related to the tribe Sciurini, based on characters of the skull, the baculum, and the number of mammae (Moore, 1959, 1961; Pocock, 1923; Wade and Gilbert, 1940). Analyses of nuclear (Steppan et al., in press) and combined nuclear and mitochondrial (Mercer and Roth, 2003) DNA, however, recovered no particularly close relationship between the Marmotini + Tamiini and the Sciurini, finding instead that the Marmotini and Tamiini belong to a clade also including the Chinese rock squirrel *Sciurotamias* and the tribes Xerini (African, North African, and long-clawed ground squirrels), Protoxerini (African palm, African giant, and sun squirrels), and Funambulini (African tree squirrels) minus *Funambulus*. Moore (1959) considered the Xerini basal to the Protoxerini (concluding that similarities between the Marmotini + Tamiini and the Xerini were due to convergence), and suggested a close relationship between the Ratufini (Asian giant squirrels) and the Nannosciurini (Oriental tree squirrels), and between the Protoxerini and the Funambulini.

Placement of the American red squirrels, genus *Tamiasciurus*, which along with *Sciurotamias* make up the Tamiasciurini, has historically been problematic. The unique genital tract of this group (Mossman et al., 1932; Pocock, 1923) led Pocock (1923) to place it in a separate tribe, the Tamiasciurini. Simpson (1945) retained this designation, but Layne (1952) argued that the vestigial baculum was insufficient to warrant tribal status for *Tamiasciurus*. Moore (1959) suggested affinities between *Tamiasciurus* and the Xerini, and a distant relationship with the Sciurini. However, close affinities with *Sciurus* have been supported by studies based on fossils, protein electrophoresis, and immunological data (Black, 1963; Bryant, 1945; Ellis and Maxson, 1980; Hafner et al., 1994; Hight et al., 1974). Analyses of nucleotide data

(Mercer and Roth, 2003; Steppan et al., in press) found *Tamiasciurus* basal among the Sciurini (minus *Sciurillus*). Mercer and Roth (2003) found *Sciurotamias* basal to a clade including the Marmotini, Tamiini, Protoxerini, and Funambulini (minus *Funambulus*), while Steppan et al. (in press) found *Sciurotamias* nested within this clade, sister to the Marmotini.

Two recent studies have provided considerable insight regarding relationships among and generic content of tribes and have presented evidence for intergeneric relationships. Mercer and Roth (2003) inferred relationships for all but one extant sciurid genus using one nuclear (IRBP; interphotoreceptor retinoid binding protein) and two mitochondrial (12S and 16S rRNA) genes. The deepest divergence within the family in their reconstruction was that of *Sciurillus*, followed by *Ratufa*, from the other sciurids. The remaining genera were resolved as three major lineages: the Pteromyinae, Tamiasciurini, and Sciurini (except *Sciurillus*); the Nannosciurini plus *Funambulus*; and the Marmotini, Tamiini, Protoxerini, and Xerini plus *Sciurotamias* and most members of the Funambulini. Steppan et al. (in press) used two nuclear genes (c-myc and RAG1) to infer relationships among 23 squirrel genera. Their results were in broad agreement with those of Mercer and Roth (2003), but differ in branching order and in their placement of *Sciurotamias*. Neither of these studies, however, was designed to investigate the specific content and monophyly of genera.

In this study, we use mitochondrial cytochrome *b* (cyt *b*) gene sequences from a large and densely sampled array of sciurid taxa to further clarify the evolutionary relationships within the Sciuridae. We aimed at complementing the taxon sampling design of Mercer and Roth (2003) and Steppan et al. (in press) by densely sampling from the intraspecific level to the tribal level. This approach allows evaluation of the monophyly of genera and, in some cases, species. Additionally, we re-examine hypotheses for higher-level relationships within the Sciuridae and the monophyly of the family. We use both maximum parsimony and Bayesian phylogenetic methods to reconstruct the phylogeny of sciurids. For Bayesian phylogenetic reconstruction we test conventional and more complex models of nucleotide substitution to identify a best fitting model for the data. Estimated phylogenies are used to evaluate the validity of the current taxonomy at the generic level, as well as for higher taxa.

2. Materials and methods

2.1. Taxon sampling

Within the family Sciuridae, our taxonomic sampling (Table 1) included representatives of 12 species in six

Table 1
Taxa used in this study

Tribe/Family/Subfamily	O.T.U.	Accession number	Sequence length (bp)	
Aplodontidae	<i>Aplodontia rufa</i>	AJ389528	1140	
Myoxidae	<i>Eliomys quercinus</i>	EQAJ5030	1140	
	<i>Glis glis</i>	AJ225031	1140	
	<i>Graphiurus murinus</i>	AJ225115	1140	
Anomaluridae	<i>Anomalurus</i> sp.	AJ389526	1140	
	<i>Idiurus macrotis</i>	AJ389525	1140	
Pedetidae	<i>Pedetes surdaster</i>	AJ389527	1140	
Castoridae	<i>Castor fiber</i>	AJ389529	1140	
Dipodidae	<i>Allactaga elater</i>	AJ389534	1140	
Caviidae	<i>Cavia porcellus</i>	NC000884	1140	
Pteromyinae	<i>Belomys pearsonii</i>	AB030262	1068	
	<i>Glaucomys sabrinus</i>	AF030390	1140	
	<i>Glaucomys sabrinus</i>	AF271668	1140	
	<i>Glaucomys sabrinus</i>	AF359211	1140	
	<i>Glaucomys sabrinus</i>	AF359221	1140	
	<i>Glaucomys sabrinus griseifrons</i>	AF359238	1140	
	<i>Glaucomys volans</i>	AJ389531	1140	
	<i>Glaucomys volans</i>	AF157921	1140	
	<i>Hylopetes phayrei</i>	AB030259	1068	
	<i>Petaurista alborufus</i>	AB023902	1068	
	<i>Petaurista elegans</i>	AB047380	1068	
	<i>Petaurista leucogenys</i>	AB023906	1068	
	<i>Petaurista petaurista</i>	AB023909	1068	
	<i>Petaurista philippensis</i>	AB023907	1068	
	<i>Petinomys setosus</i>	AB030260	1068	
	<i>Pteromys momonga</i>	AB030263	1068	
	<i>Pteromys volans</i>	AB023910	1068	
	Funambulini	<i>Paraxerus cepapi</i>	PCU59179	714
	Marmotini	<i>Ammospermophilus harrisi</i>	AF157926	1140
		<i>Ammospermophilus harrisi</i>	AHU46172	528
<i>Ammospermophilus interpres</i>		AIU46174	528	
<i>Ammospermophilus leucurus</i>		ALU46175	528	
<i>Cynomys gunnisoni</i>		AF157930	1140	
<i>Cynomys leucurus</i>		AF157879	1140	
<i>Cynomys ludovicianus</i>		AF157890	1140	
<i>Cynomys mexicanus</i>		AF157847	1140	
<i>Cynomys parvidens</i>		AF157922	1140	
<i>Marmota baibacina centralis</i>		AF143915	1140	
<i>Marmota baibacina kastchenkoi</i>		AF143914	1140	
<i>Marmota bobak bobak</i>		AF143916	1140	
<i>Marmota bobak bobak</i>		AF143917	1140	
<i>Marmota broweri</i>		AF143919	1140	
<i>Marmota caligata caligata</i>		AF143920	1140	
<i>Marmota camtschatica camtschatica</i>		AF143921	1140	
<i>Marmota camtschatica doppelmayri</i>		AF143922	1140	
<i>Marmota caudata aurea</i>		AF143925	1140	
<i>Marmota caudata caudata</i>		AF143923	1140	
<i>Marmota flaviventris luteola</i>		AF143926	1140	
<i>Marmota flaviventris obscura</i>		AF143927	1140	
<i>Marmota himalayana robusta</i>		AF143928	1140	
<i>Marmota marmota</i>		AF100711	1140	
<i>Marmota marmota marmota</i>		AF143930	1140	
<i>Marmota marmota marmota</i>		AF143929	1140	
<i>Marmota menzbieri zachidovi</i>		AF143931	1140	
<i>Marmota monax monax</i>		AF143934	1140	
<i>Marmota monax ochracea</i>		AF143932	1140	
<i>Marmota olympus</i>		AF111182	810	
<i>Marmota sibirica sibirica</i>		AF143937	1140	
<i>Marmota vancouverensis</i>		AF143939	1140	
<i>Spermophilus adocetus</i>		AF157844	1140	
<i>Spermophilus annulatus</i>	AF157851	1140		
<i>Spermophilus armatus</i>	AF157901	1140		
<i>Spermophilus atricapillus</i>	AF157945	1140		

Table 1 (continued)

Tribe/Family/Subfamily	O.T.U.	Accession number	Sequence length (bp)
	<i>Spermophilus beechyi</i>	AF157918	1140
	<i>Spermophilus beldingi creber</i>	AF157951	1140
	<i>Spermophilus brunneus brunneus</i>	AF157952	1140
	<i>Spermophilus citellus</i>	AF157859	1140
	<i>Spermophilus columbianus columbianus</i>	AF157939	1140
	<i>Spermophilus columbianus ruficaudus</i>	AF157943	1140
	<i>Spermophilus dauricus</i>	AF157899	1140
	<i>Spermophilus elegans elegans</i>	AF157891	1140
	<i>Spermophilus erythrogegens</i>	AF157855	1140
	<i>Spermophilus erythrogegens</i>	AF157875	1140
	<i>Spermophilus franklinii</i>	AF157893	1140
	<i>Spermophilus franklinii</i>	AF157894	1140
	<i>Spermophilus fulvus</i>	AF157913	1140
	<i>Spermophilus lateralis trepidus</i>	AF157887	1140
	<i>Spermophilus lateralis trepidus</i>	AF157950	1140
	<i>Spermophilus madrensis</i>	AF157946	1140
	<i>Spermophilus madrensis</i>	AF157947	1140
	<i>Spermophilus major</i>	AF157903	1140
	<i>Spermophilus major</i>	AF157905	1140
	<i>Spermophilus mexicanus mexicanus</i>	AF157848	1140
	<i>Spermophilus mexicanus parvidens</i>	AF157853	1140
	<i>Spermophilus mojavensis</i>	AF157928	1140
	<i>Spermophilus mollis</i>	AF157938	1140
	<i>Spermophilus musicus</i>	AF157904	1140
	<i>Spermophilus pallicaudata</i>	AF157869	1140
	<i>Spermophilus parryi</i>	AF157896	1140
	<i>Spermophilus parryi kenicotti</i>	AF157931	1140
	<i>Spermophilus parryi plesius</i>	AF157927	1140
	<i>Spermophilus perotensis</i>	AF157948	1140
	<i>Spermophilus pygmaeus</i>	AF157907	1140
	<i>Spermophilus pygmaeus</i>	AF157910	1140
	<i>Spermophilus relictus</i>	AF157862	1140
	<i>Spermophilus relictus</i>	AF157867	1140
	<i>Spermophilus relictus</i>	AF157876	1140
	<i>Spermophilus richardsoni</i>	AF157915	1140
	<i>Spermophilus saturatus</i>	AF157917	1140
	<i>Spermophilus spilosoma</i>	AF157846	1140
	<i>Spermophilus spilosoma marginatus</i>	AF157911	1140
	<i>Spermophilus suslicus</i>	AF157895	1140
	<i>Spermophilus suslicus</i>	AF157898	1140
	<i>Spermophilus tereticaudus</i>	AF157941	1140
	<i>Spermophilus townsendii idahoensis</i>	AF157949	1140
	<i>Spermophilus townsendii townsendii</i>	AF157935	1140
	<i>Spermophilus townsendii vigilis</i>	AF157889	1140
	<i>Spermophilus tridecemlineatus arenicola</i>	AF157877	1140
	<i>Spermophilus undulatus</i>	AF157906	1140
	<i>Spermophilus undulatus</i>	AF157912	1140
	<i>Spermophilus variegatus</i>	AF157854	1140
	<i>Spermophilus variegatus utah</i>	AF157878	1140
	<i>Spermophilus washingtoni</i>	AF157936	1140
	<i>Spermophilus xanthoprimum</i>	AF157909	1140
Nannosciurini	<i>Callosciurus erythraeus</i>	AB043876	1080
	<i>Callosciurus erythraeus</i>	AB043877	1080
	<i>Callosciurus finlaysonii</i>	AB043878	1080
	<i>Callosciurus nigrovittatus</i>	AB043882	1080
	<i>Callosciurus prevostii</i>	AB043879	1080
	<i>Callosciurus prevostii</i>	AB043880	1080
Sciurini	<i>Microsciurus flaviventer</i>	MFU46169	528
	<i>Sciurillus pusillus</i>	SPU46179	528
	<i>Sciurus aberti barberi</i>	SAU10177	1134
	<i>Sciurus aberti chuscensis</i>	SAU10176	1134
	<i>Sciurus aberti kaibabensis</i>	SAU10182	1140
	<i>Sciurus aestuans</i>	SAE389530	1140
	<i>Sciurus carolinensis</i>	SCU46167	1140

Table 1 (continued)

Tribe/Family/Subfamily	O.T.U.	Accession number	Sequence length (bp)
Tamiasciurini	<i>Sciurus lis</i>	AB030024	1040
	<i>Sciurus lis</i>	AB043881	1080
	<i>Sciurus niger</i>	SNU10180	1140
	<i>Sciurus stramineus</i>	AB030025	1040
	<i>Sciurus vulgaris</i>	AB030028	1040
	<i>Sciurus vulgaris</i>	AJ238588	1140
	<i>Tamiasciurus douglasii</i>	AF322953	402
	<i>Tamiasciurus hudsonicus</i>	AB030029	1040
	<i>Tamiasciurus mearnsi</i>	AF322954	402
	Tamiini	<i>Eutamias sibiricus</i>	AF147666
<i>Neotamias amoenus</i>		AF147630	1140
<i>Neotamias amoenus</i>		AF147632	1140
<i>Neotamias bulleri</i>		AF147634	1140
<i>Neotamias canipes</i>		AF147635	1140
<i>Neotamias cinericollis</i>		AF147636	1140
<i>Neotamias cinericollis</i>		AF147638	1133
<i>Neotamias dorsalis</i>		AF147639	1133
<i>Neotamias dorsalis</i>		AF147641	1138
<i>Neotamias durangae</i>		AF147642	1140
<i>Neotamias merriami</i>		AF147644	1140
<i>Neotamias minimus</i>		AF147645	1131
<i>Neotamias minimus</i>		AF147650	1140
<i>Neotamias obscurus</i>		AF147651	1139
<i>Neotamias ochrogenys</i>		AF147654	1136
<i>Neotamias palmeri</i>		AF147655	1103
<i>Neotamias panamintinus</i>		AF147656	1140
<i>Neotamias quadrimaculatus</i>		AF147657	1140
<i>Neotamias quadrivittatus</i>		AF147658	1140
<i>Neotamias quadrivittatus</i>		AF147660	1140
<i>Neotamias ruficaudus</i>		AF147661	1140
<i>Neotamias rufus</i>		AF147662	1140
<i>Neotamias rufus</i>		AF147663	1140
<i>Neotamias senex</i>		AF147665	1140
<i>Neotamias siskiyou</i>		AF147668	1140
<i>Neotamias sonomae</i>		AF147669	1118
<i>Neotamias townsendii</i>		AF147674	1140
<i>Neotamias townsendii</i>		AF147676	1140
<i>Neotamias umbrinus</i>		AF147677	1070
<i>Tamias striatus</i>	AF147673	1139	
Xerini	<i>Spermophilopsis leptodactylus</i>	AF157865	1140
	<i>Xerus inauris</i>	AY452689	1140
	<i>Xerus rutilus</i>	AY452690	1140

genera of the subfamily Pteromyinae and 102 species in 15 genera of the subfamily Sciurinae. Two or more individuals were included for 37 species, with multiple nominal subspecies used where available. Multiple, successive outgroups (Smith, 1994) were used to root phylogenies and to examine the monophyly of the Sciuridae with respect to the Aplodontidae, the hypothesized sister family to the Sciuridae (Adkins et al., 2001; DeBry and Sagel, 2001; Huchon et al., 1999, 2000; Lavocat and Parent, 1985; Montgelard et al., 2002; Nedbal et al., 1996; Stepan et al., in press; Wahlert, 1985). Aplodontidae contains only a single extant representative, the mountain beaver (*Aplodontia rufa*).

In addition to *Aplodontia* (left as a member of the ingroup), outgroup taxa from other rodent families included two members of the Anomaluridae (*Idiurus macrotis* and *Anomalurus* sp.), one member of the

Castoridae (*Castor fiber*), one member of the Dipodidae (*Allactaga elater*), three members of the Gliridae (*Eliomys quercinus*, *Glis glis* and *Graphiurus murinus*), one member of the Pedetidae (*Pedetes surdaster*), and a hystricomorph rodent (*Cavia porcellus*). The majority of sequences used in this study (167 of 169 total sequences) have been previously published (Table 1). Most of these sequences were originally published in genus- or species-level phylogenies (e.g., Arbogast et al., 2001; Oshida et al., 2000; Oshida and Masuda, 2000; Oshida et al., 2001; Piaggio and Spicer, 2001; Stepan et al., 1999; Wettstein et al., 1995) or as direct submissions to GenBank. This study is the first time these sequences have been combined in one analysis to explore higher-level relationships. This paper provides two new complete cyt *b* sequences (*Xerus inauris* and *X. rutilus*), which have been deposited in GenBank (see Table 1).

2.2. DNA sequencing and sequence alignment

Total DNA was extracted from 95% ethanol-preserved liver samples from *X. inauris* and *X. rutilus* using standard proteinase K digestion followed by phenol/chloroform/isoamyl alcohol organic separation methods as described in Sambrook and Russell (2001). The *cyt b* gene was PCR-amplified (Saiki et al., 1988) using the primers L14725 (Irwin et al., 1991) and XerusREV (5'-TTNGGTTTCAAGACCAAGT-3', this study), located in the flanking tRNA-Glutamine and tRNA-Threonine genes. PCR amplification was carried out using an initial denaturation at 94 °C for 3 min 45 s; 45 cycles of denaturation at 94 °C for 1 min, annealing at 45 °C for 1 min, and extension at 72 °C for 2 min; and a final extension at 72 °C for 7 min. Products from PCR were excised from 1.4% agarose electrophoresis gels and purified using the MinElute Gel Extraction Kit (Qiagen). Purified PCR products were sequenced in both directions using the amplification primers and the internal primer XerusFctb (5'-TGAGGRCAAATATCCTTCTGAGG-3', this study) with the CEQ Dye Terminator Cycle Sequencing Quick Start Kit (Beckman Coulter) and run on a Beckman CEQ2000 automated sequencer according to the manufacturer's protocols. Raw sequence chromatographs from novel sequences were edited in Sequencher 3.1 (Gene Codes).

All sequences were aligned manually in GeneDoc 2.6.002 (Nicholas and Nicholas, 1997). Aligned DNA sequences were translated to amino acid sequences to confirm alignment and check for the presence of stop codons or open reading frame shifts (neither of which were detected), which might have indicated pseudogenes. Since the *cyt b* gene is protein coding, indels were infrequent and occurred only in units representing the gain or loss of complete codons. Only two indels were observed, both 3 bp in length and both inferred as deletions in *Sciurus aberti barberi* and *S. a. chuscensis*. The exact position of one of these, beginning at position 19 or 22, could not be confidently determined, so three nucleotides were conservatively coded as missing data for these taxa. The second inferred deletion occurred at positions 694–696, and alignment was otherwise unambiguous. The alignment (in Nexus format) is available at TreeBASE (<http://treebase.bio.buffalo.edu/treebase>) under the matrix Accession No. M1630.

2.3. Phylogenetic reconstruction

Phylogenetic relationships were estimated using both maximum parsimony (MP) and Bayesian Metropolis-coupled Markov Chain Monte Carlo phylogenetic methods (MCMC). Unweighted MP searches were conducted in PAUP* v4.0b10 (Swofford, 2002) with gaps treated as missing. The heuristic search strategy was used to search for optimal MP trees with 100 random taxon addition sequence replicates and starting

trees obtained via stepwise addition. Settings for MP analyses included tree bisection–reconnection branch swapping, steepest descent off, and MULTREES option on (Swofford, 2002). Clade support was assessed with 300 bootstrap pseudoreplicates, each with 10 random addition sequence replicates. Additional parameters employed for the bootstrap analysis were identical to those described above.

All MCMC phylogenetic reconstructions were conducted using MrBayes 2.10 (Huelsenbeck and Ronquist, 2001) with uniform priors and model parameters estimated as part of the analyses. Three heated chains and a single cold chain were used in all MCMC analyses, and runs were initiated with random trees. Trees were sampled every 100 generations, and majority rule consensus phylograms and posterior probabilities for nodes were assembled from all post-burn-in sampled trees. The data were subjected to MCMC analyses under multiple evolutionary models that differed in the way they parameterized among-site rate variation. Under each model of evolution, four independent MCMC runs were conducted, each with 1.4 million generations per run.

We used ModelTest 3.0 (Posada and Crandall, 1998) to infer the simplest, best-fit model of evolution based on hierarchical log-likelihood ratio tests comparing successively more complex models (Huelsenbeck and Crandall, 1997; Posada and Crandall, 2001). The most complex (parameter rich) model that ModelTest 3.0 can evaluate is a General Time Reversible (GTR, Tavaré, 1986) model with a parameter (γ) estimating among-site rate variation (Yang, 1993) and another parameter (I) describing the proportion of invariant sites. This GTR + G + I model was selected by ModelTest, but we also wished to explore the effects of more complex models on topology and optimized tree likelihood scores (used to represent model fit to data).

MrBayes is capable of employing more complex models than this GTR + G + I model. In particular, MrBayes allows among-site rate variation to be partitioned among defined sites (site-specific γ) as well as allowing the use of an autocorrelation parameter to account for autocorrelation of rates among sites (Kimura, 1986; Nielsen, 1997; Schöniger and Von Haeseler, 1994; Yang, 1995). These two types of modifications to simple parameterization of among-site rate variation using a single γ shape parameter may be used independently as well as simultaneously in a given model. We tested the performance of four alternative models, all employing a single GTR substitution matrix, which differed only with respect to how among site rate variation was parameterized, including: GTR + G + I (chosen by Modeltest), GTR + AG (GTR model with autocorrelated among-site rate variation), GTR + partG (GTR model with site-specific partitions of γ , partitioned as described below), and GTR + partAG (GTR with site-specific partitions of γ with autocorrelated rates).

The structure of the *cyt b* protein has been well characterized (Howell, 1989), and variation of evolutionary rates across functional domains has been analyzed for birds and mammals (Griffiths, 1997; Irwin et al., 1991). For the site-specific gamma parameter partition models (GTR + partG and GTR + partAG) we assigned sites to one of three partitions, according to the structural model of the *cyt b* gene presented by Griffiths (1997) for the Falconidae (Aves). Because the *cyt b* gene is 1143 bp long in Falconidae and only 1140 bp in most sciurids, an alignment was created in Genedoc (Nicholas and Nicholas, 1997) including several bird and squirrel reference sequences. A single 3 bp indel at positions 4–6 accounted for the difference in length, and so we divided the squirrel gene into the following partitions: intermembrane (1–90, 316–336, 613–678, 925–954, and 1111–1140), transmembrane (91–165, 238–315, 337–399, 532–612, 679–744, 856–924, 955–1023, and 1051–1110), and matrix (166–237, 400–531, 745–855, and 1024–1050).

To explore whether these three partitions in fact appear to evolve under different patterns of among-site rate variation (justifying a model to account for this), we conducted MCMC runs in MrBayes independently on each partition. These runs employed a GTR + G model and were run in triplicate to guard against any one run failing to reach a global optimum of likelihood stationarity. Details of MCMC analyses were the same as those described previously.

2.4. Hypothesis testing

We examine two types of hypotheses with alternative Bayesian MCMC analyses: (1) hypotheses regarding which model of evolution best fits the data (alternative models of evolution explored are discussed above) and (2) null hypotheses of monophyly of taxa that were recovered as paraphyletic in the unconstrained MCMC reconstruction. To test alternate topological hypotheses, we conducted independent MCMC analyses in which a particular group was constrained as monophyletic based on the current taxonomy as described by McKenna and Bell (1997). To evaluate both types of hypotheses, we compared the 95% credibility interval (CI) of MCMC chain likelihood scores between the null and alternative hypotheses. To calculate the 95% CI, we ranked all post-burn-in tree estimates by ln likelihood and included the most likely 95% (Felsenstein, 1968; Huelsenbeck et al., 2002). Since a 95% Bayesian credibility interval is similar to a 95% confidence interval of classical statistics (Huelsenbeck et al., 2002), the degree of overlap observed between null and alternative ln likelihood CIs was used to test hypotheses. In testing null hypotheses based on current taxonomy, the null constrained hypotheses were rejected if the 95% CI of constrained MCMC chain likelihood scores did not overlap with the 95% CI of unconstrained MCMC chain likelihood scores.

3. Results

3.1. Maximum parsimony phylogenetic analysis

Of the 1140 aligned bases (380 codons), 446 were invariant (iv), 97 variable characters were parsimony uninformative (vpu), and 597 were parsimony informative (pi). Among the three codon positions, the first position contained 197 iv, 37 vpu, and 146 pi characters; the second position contained 247 iv, 58 vpu, and 75 pi characters; and the third position contained 2 iv, 2 vpu, and 376 pi characters. The observed base composition for the L-strand was: A = 28.42%; C = 27.07%; G = 12.63%; and T = 31.88%. Uncorrected percent pairwise sequence divergence was highest between *C. porcellus* and *P. surdaster* (27.54%). Within the Sciuridae, the highest observed uncorrected pairwise divergence was between *Spermophilus dauricus* and *Paraxerus cepapi* (25.82%).

The unweighted MP reconstruction resulted in 208 equally parsimonious trees with length = 9289 steps, consistency index = 0.137, retention index = 0.667, re-scaled consistency index = 0.091, and homoplasy index = 0.863. A strict consensus of all 208 optimal MP trees is shown in Fig. 1, with bootstrap percentages (BP) above 50% indicated. Three major clades are evident among sciurids. The first is a weakly supported clade including *Aplodontia*, *Sciurillus*, and *Paraxerus* as sister to a monophyletic *Callosciurus* (BP = 99%). The second includes a weakly supported tribe Xerini, the Pteromyinae, the Tamiasciurini (BP = 100%), and the Sciurini minus *Sciurillus* (BP = 92%). Monophyly of the genera *Xerus* (BP = 100%), *Petaurista* (BP = 94%), *Glaucomys* (BP = 100%), *Pteromys* (BP = 100%), and *Tamiasciurus* (BP = 100%) received strong support. *Sciurus* was resolved as paraphyletic with respect to *Microsciurus*. The third clade shows the tribe Tamiini sister to the Marmotini (BP = 76%).

Among the species for which two or more individuals were included, the following were not resolved as exclusive monophyletic groups (i.e., individuals of the same species failed to group together) in the MP analysis: *Glaucomys sabrinus*, *Neotamias cinericollis*, *N. dorsalis*, *N. minimus*, *Ammospermophilus harrisi*, *Spermophilus citellus*, *S. major*, *S. mexicanus*, *S. pygmaeus*, *S. spilosoma*, and *S. townsendi*. *Neotamias rufus* was part of an unresolved polytomy with *N. quadrivittatus* and a clade including *N. cinericollis*, *N. dorsalis*, *N. palmeri*, and *N. umbrinus*.

3.2. Bayesian phylogenetic analyses

Based on hierarchical log-likelihood ratio tests of successively more complex models of sequence evolution, Modeltest indicated that the simplest, best-fit model for the dataset was the GTR + G + I model with the following




Fig. 1. Strict consensus of all 208 equally optimal trees based on unweighted maximum parsimony search. Bootstrap percentages above 50% are provided. Higher taxa are also given. Seven clades representing independent lineages containing members of the genus *Spermophilus* are indicated by sequentially labeled shading (S-1 to S-7).

parameter estimates: [A–C] = 0.209, [A–G] = 10.666, [A–T] = 0.413, [C–G] = 0.752, [C–T] = 4.817, [G–T] = 1.000, gamma parameter [G] = 0.5826, proportion of invariant sites [I] = 0.342.

Independent MCMC on the defined partitions (to initially detect domain-specific patterns of among-site rate variation) all converged to a common stationary

value across replicate runs. Estimates of the gamma shape parameter for each domain did show domain-specific differentiation in the mean values (but overlap of 95% credibility interval) for gamma parameter estimates as follows (gamma shape parameter estimates reported as means with the minimum and maximum estimated values within the 95% credibility interval of trees):

intermembrane partition = 0.3197 (0.2222–0.4292), transmembrane partition = 0.3279 (0.2730–0.3804), matrix partition = 0.3030 (0.2459–0.3638).

The pooled mean ln likelihood score for the post-burn-in trees recovered by the four MCMC runs using the GTR + AG model was $-37,999.54$ (95% credibility interval $-38,030.14$ to $-37,970.18$). That of the GTR + partG model was $-38,037.56$ ($-38,068.62$ to $-38,008.20$), and that of the GTR + partAG model was $-37,997.51$ ($-38,036.15$ to $-37,967.36$) (Fig. 2). These three models have equal (GTR + AG) or greater numbers of model parameters as the GTR + G + I model. Despite being equally or more complex models, all these alternative models resulted in lower MCMC chain likelihood score values than did MCMC chains employing the GTR + G + I model (mean $-37,934.99$; 95% credibility interval $-37,870.63$ to $-37,960.82$; Fig. 2). Because these equally (GTR + AG) or more complex models yielded less optimal likelihood scores for chains than the GTR + G + I model, we chose to report our conclusions based on the latter (Fig. 3). The burn-in plots of the four independent MCMC runs using the GTR + G + I model (Fig. 4) show that all four runs reached stationarity prior to 350 000 generations. Thus, our burn-in time of 400 000 generations was sufficient. All four runs of the GTR + G + I model converged with regard to chain likelihood scores (Fig. 4), topology (50% majority rule consensus identical among runs), and

posterior probability estimates for clade support. Also, parameter plots for all four independent runs show overlapping estimates of substitution model parameters at stationarity (Fig. 5). Parameter estimates based on the combination of all post-burn-in generations from the four independent MCMC runs under the GTR + G + I model, reported as means with the minimum and maximum estimated values within the 95% credibility interval of trees, were: $[A-C] = 0.242$ (0.529–0.146), $[A-G] = 13.171$ (21.265–8.455), $[A-T] = 0.479$ (0.910–0.234), $[C-G] = 0.864$ (1.797–0.338), $[C-T] = 5.851$ (11.812–3.443), $[G-T] = 1$, gamma parameter $[G] = 0.584$ (0.672–0.499), and proportion of invariant sites $[I] = 0.294$ (0.347–0.237).

Based on the evidence for convergence among runs, we pooled all post-burn-in trees (representing a total of 4 million post-burn-in generations) to estimate the posterior probabilities of clades. A 50% majority rule consensus phylogram (branch lengths averaged over all post-burn-in trees) is presented in Fig. 3 with posterior probabilities from all post-burn-in generations of the GTR + G + I model runs indicated. *Aplodontia* appears sister to the Sciuridae with 99% Bayesian posterior probability (PP). The earliest divergence within the Sciuridae is that of *Sciurillus*, followed by the Xerini, *Paraxerus*, and *Callosciurus*. Among the remaining squirrels, three major clades are apparent: the *Tamiasciurini*, *Sciurini* (minus *Sciurillus*), and *Pteromyinae* (PP = 98%); the *Tamiini* (PP = 99%); and the *Marmotini* (PP = 100%). Monophyly of flying squirrels is strongly supported (PP = 99%), as is that of the genera *Ammodontomys* (PP = 90%), *Callosciurus* (PP = 100%), *Cynomys* (PP = 100%), *Marmota* (PP = 100%), *Sciurus* (PP = 100% including *Microsciurus*), *Neotamias* (PP = 100%), and *Tamiasciurus* (PP = 100%). *Microsciurus* is deeply nested within the *Sciurus* clade.

As in the MP reconstruction, the MCMC analysis recovered *Spermophilus (Ictidomys) mexicanus* and *S. (I.) tridecemlineatus* as sister species (PP = 100%, BP = 100%) basal to the clade (PP = 93%, BP < 50%) containing *S. (Poliocitellus) franklinii*, *Cynomys*, *S. (Xerospermophilus) mohavensis*, *S. (X.) tereticaudus*, and the remaining members of the subgenus *Ictidomys*, *S. (I.) perotensis* and *S. (I.) spilosoma*. The remaining members of subgenus *Ictidomys* are sister to subgenus *Xerospermophilus* (PP = 100%, BP = 87%). The subgenus *Spermophilus* forms two distinct clades, one consisting entirely of Old World taxa (PP = 100%, BP = 92%) and one of New World taxa plus *S. undulatus* (PP = 99%, BP = 90%). As in the MP reconstruction, multiple members of the same species failed to group together (i.e., the same species did not appear monophyletic) as in the MCMC analysis, with the addition of *N. rufus*.

Based on the resolution of clades from the GTR + G + I model, we identified seven relationships




Fig. 2. Comparison of MCMC chain likelihood scores across alternative models of evolution evaluated. GTR + G + I = General Time Reversible (GTR) with a gamma distribution of among-site rate variation and a proportion of sites invariant; GTR + AG = GTR with autocorrelated gamma; GTR + partG = GTR with site-specific partitions of gamma; GTR + partAG = GTR with site-specific partitions of gamma with autocorrelated rates. Likelihood scores for each model are reported as the mean (box) and 95% credibility interval (bars) based on the combined distribution of 4 million post-burn-in generations (from four independent MCMC runs per model).




Fig. 3. Majority rule consensus phylogram (with branch lengths averaged over all trees) resulting from Bayesian phylogenetic analysis using a GTR + G + I model of evolution. Posterior probabilities estimated from a total of 4 million post-burn-in generations (from four independent runs) are indicated for major nodes. Higher taxon names are also given. Seven clades representing independent lineages containing members of the genus *Spermophilus* are indicated by sequentially labeled shading (S-1 to S-7).




Fig. 4. Burn-in plot of ln likelihood scores of MCMC chains across 1.4 million generations for four independent runs under the GTR + G + I model of evolution.




Fig. 5. Plots of selected parameters of independent MCMC runs under the GTR + G + I model through 1.4 million generations. All four independent runs are plotted per graph to show common burn-in rates and similar, overlapping parameter estimates. (A) Plot of the parametric estimates of $r(A-G)$ from the GTR rate matrix. (B) Plot of the parametric estimates of $r(A-T)$ from the GTR rate matrix. (C) Plot of the gamma parameter (Γ) estimates. (D) Plot of the proportion of invariant sites parameter (I) estimates.

that contradicted the null hypothesis of monophyly of conspicuous taxonomic groups. These included the paraphyly of the tribe Sciurini, the genera *Sciurus* and *Spermophilus*, and the subgenera *Ictidomys*, *Spermophilus*, and *Otospermophilus*. Finally, the Old World species

Spermophilus undulatus is nested within the clade containing the New World members of the subgenus *Spermophilus*. A comparison between MCMC chain likelihood scores of the unconstrained GTR + G + I runs and MCMC runs where these particular relationships




Fig. 6. Comparison between the ln likelihood score for MCMC chains for the unconstrained GTR + G + I model runs and those of alternative runs (with the same model) where particular relationships were constrained. Names of constrained runs represent the groups restricted to monophyly. Likelihood scores are reported as the mean (box) and 95% credibility interval (bars). The shaded area represents the likelihood score range spanned by the 95% credibility interval of the unconstrained MCMC analyses, and represents the region where null hypotheses of constrained relationships would not be rejected.

were constrained is provided in Fig. 6. The MCMC searches constrained to monophyly of the genera *Sciurus* and *Spermophilus* and of the *Spermophilus* subgenera *Spermophilus* and *Otospermophilus* resulted in chain likelihood scores with 95% CIs below the lower bound of that of the unconstrained runs (Fig. 6). Given these results, we reject the null hypothesis of monophyly for of each of these groups. Although we could not reject monophyly of the tribe Sciurini, the *Spermophilus* subgenus *Ictidomys*, and the Old World members of the subgenus *Spermophilus*, these groups did not appear monophyletic in either the MP or the unconstrained MCMC analyses.

4. Discussion

4.1. Models of evolution

Our evaluation of alternative complex models of nucleotide substitution for reconstructing phylogenies with Bayesian MCMC methods yielded unexpected, interesting results. Previous authors have estimated rates and patterns of nucleotide substitution for different domain-encoding regions of the *cyt b* gene, and have found domain-specific rates of substitution (Griffiths, 1997; Irwin et al., 1991). Our initial exploratory independent MCMC estimation of the among-site rate parameter (γ) for each of the three partitions did provide evidence for at least weakly differentiated domain-specific patterns of among-site rate variation. We used this preliminary evidence as justification for testing

the GTR + partG and GTR + partAG models that attempt to rescale a general gamma shape parameter to adjust fit across the three partitions. We expected this to yield models of nucleotide evolution that more realistically accounted for patterns of nucleotide substitution by allowing partition-specific scaling of the gamma shape parameter. Instead, we found that these structurally partitioned models (GTR + partG and GTR + partAG) were a poorer fit to the data than the standard GTR + G + I model. Krajewski and King (1996) found that although divergence among cytochrome *b* regions differed by as much as fivefold, variation was not correlated with structural domains. Their findings, together with the results of our model evaluation, suggest that while certain characteristics of molecular evolution are almost certainly related to the structure of the protein encoded by a gene, patterns of among-site rate variation are do not appear specific to particular domains or groups of domains (at least within the *cyt b* gene in sciurids). In addition to partitioned gamma models, the GTR + G + I model outperformed the GTR + AG model. These results seem to support the strong utility of the GTR + G + I model and, in particular, the ability of a simple gamma parameter to explain among-site rate variation when the proportion of invariant sites is estimated.

4.2. Relationships and taxonomy of the Sciuridae

A sister-group relationship between the Sciuridae and the Aplodontidae is supported by analyses of mitochondrial (Montgelard et al., 2002; Nedbal et al., 1996) and nuclear (Adkins et al., 2001; DeBry and Sagel, 2001; Huchon et al., 1999, 2000) gene sequences as well as by morphological data (Lavocat and Parent, 1985; Wahlert, 1985). Most recently, Mercer and Roth (2003) and Steppan et al. (in press) also provided strong evidence for Aplodontidae being the sister taxon to a monophyletic Sciuridae. While our MP reconstruction did not recover this relationship, our MCMC phylogeny did support this relationship between families, but failed to find strong support (PP = 41%) for the monophyly of the Sciuridae. Because our analyses do not contradict those of Mercer and Roth (2003) and Steppan et al. (in press), both of which employed more slowly evolving nuclear genes, we consider a sister-group relationship between Aplodontidae and a monophyletic Sciuridae to be the favored hypothesis.

In interpreting the results of this study, it is important to consider taxonomic sampling biases and how they may affect perspectives regarding the major relationships among sciurids. Our sampling includes six of the 16 extant genera within the Pteromyinae, and 15 of the 37 extant genera in seven of the nine recognized tribes within the Sciurinae (McKenna and Bell, 1997). Among the sciurine tribes, our generic taxon sampling lacks the

Protoxerini and the monotypic Ratufini and is deficient within the tribes Funambulini (one of four) and Nannosciurini (one of thirteen). Our generic sampling is, however, complete for the tribes Marmotini, Tamiasciurini, and Tamiini and nearly complete for the Xerini (two of three) and Sciurini (three of four). The basal position of the Ratufini, along with the diversity of the Nannosciurini suggested by Mercer and Roth (2003) and Steppan et al. (in press), implies that our taxon sampling may limit conclusions regarding the deepest relationships among sciurids.

Subfamily classification within the family Sciuridae has been a major taxonomic problem. Several studies have suggested that the Sciurinae are paraphyletic with respect to the only other recognized subfamily, the Pteromyinae (Black, 1963; Engesser, 1979; Hight et al., 1974; James, 1963; Mein, 1970). The analyses of Mercer and Roth (2003) and Steppan et al. (in press) supported monophyly of the flying squirrels, and found them to be sister to the tribes Sciurini (minus *Sciurillus*) and Tamiasciurini. Our MCMC analysis supports these findings, showing a well-supported clade of flying squirrels (99% BPP) and a sister-group relationship with Sciurini + Tamiasciurini (98% BPP). The MP analysis shows a monophyletic Sciurini + Tamiasciurini + flying squirrel clade (<50% BS), but places *Pteromys* sister to Sciurini + Tamiasciurini (<50% BS). The present study, along with those of Mercer and Roth (2003) and Steppan et al. (in press), suggests that retention of Pteromyinae as a subfamily is not justified unless the Sciurinae is subdivided into multiple subfamilies.

Current subfamily-level designations are clearly in need of revision. Steppan et al. (in press) propose promotion of each of the five major clades recovered in their analysis to subfamily rank. The recovery of the same five clades by Mercer and Roth's (2003) analysis suggests that this reorganization is justified. However, our results demonstrate two major disagreements with the proposed classification. First, the analyses of Mercer and Roth (2003) and Steppan et al. (in press) show the tribe Xerini within the lineage (clade IV in Mercer and Roth; clade E in Steppan et al.) including the Tamiini, Marmotini, Protoxerini, and Funambulini (minus *Funambulus*). Our results suggest a much earlier divergence (more basal placement) of the Xerini. Second, their analyses show *Paraxerus* within this same clade, while our analyses instead suggest a more basal divergence of *Paraxerus*. Based on phylogenetic hypotheses presented here, together with those of Mercer and Roth (2003) and Steppan et al. (in press), the classification proposed by Steppan et al. (in press) is at least a positive step in the direction of a phylogenetic subfamilial taxonomy for the Sciuridae. However, discrepancies among all three recent phylogenetic estimates for the Sciuridae merit further verification before a stable taxonomic solution is reached. Future phylogenetic studies with increased taxonomic representation

among the basal taxa will likely be required to fully resolve this issue. Presently, a comprehensive assessment of the content of (and relationships among) the newly defined subfamilies is a major goal for Sciurid systematics. Included in this goal, efforts focused on the definition of synapomorphies supporting (or rejecting) the newly erected subfamilies are paramount.

The phylogenetic position of the monotypic genus *Sciurillus*, the Neotropical pygmy squirrel, differs between our two analyses, with neither agreeing with the current taxonomy. Based on characters of the baculum (Anthony and Tate, 1935) and skull (Moore, 1959), *Sciurillus* has been placed within the tribe Sciurini, but in its own subtribe, the Sciurillina (McKenna and Bell, 1997; Moore, 1959). This genus is unique in terms of skull characters (Moore, 1959) and jaw musculature (Ball and Roth, 1995; Thorington and Darrow, 1996), and Moore (1959) noted that it is "...one of the most distinctive genera of squirrels in the world." Both of our analyses place *Sciurillus* well outside of the remaining members of the tribe Sciurini (*Sciurus* and *Microsciurus*). In the MCMC analysis, *Sciurillus* is sister to the remaining Sciuridae (PP=41%), the monophyly of which is well supported (PP=93%). In the MP analysis, *Sciurillus* appears sister to *Aplodontia* (BP<50%). The combination of molecular evidence and morphological distinctiveness suggest that assignment of *Sciurillus* to the tribe Sciurini may be in error, and *Sciurillus* may in fact represent a basal lineage among the Sciuridae, as suggested by Mercer and Roth (2003) and Steppan et al. (in press). Unfortunately, only a partial cyt *b* sequence (528 bp) was available for this taxon, making the confidence of this diagnosis tentative.

The proposed close relationship between *Microsciurus*, the dwarf squirrel, and *Sciurus* (Hafner et al., 1994; Mercer and Roth, 2003; Steppan et al., in press) is confirmed by both analyses. Moore (1959) considered *Microsciurus* a member of the subtribe Microsciurina, along with *Leptosciurus*, *Simosciurus*, *Syntheosciurus*, and *Mesosciurus*. Because no sequence data from these genera were included in this study, the validity of their status as a subtribe cannot be evaluated, but *Microsciurus* clearly falls within a well-supported, otherwise monophyletic *Sciurus* (PP=100%, BP=92%). The designation of *Microsciurus* as a genus distinct from *Sciurus* clearly requires further attention including additional sampling of taxa within the genus *Microsciurus*.

Our analyses confirm the close relationship between *Tamiasciurus* and the Sciurini suggested by Mercer and Roth (2003) and Steppan et al. (in press). *Tamiasciurus* appears sister to *Sciurus* (including *Microsciurus*) with strong support (PP=100%, BP=78%). Either retention at the tribal level of the Tamiasciurini or inclusion of *Tamiasciurus* within the Sciurini appears justified. In either case, the analyses of Mercer and Roth (2003) and Steppan et al. (in press) show that *Sciurotamias* does not

share a close enough relationship to *Tamiasciurus* to justify its retention in the same tribe.

Piaggio and Spicer (2000, 2001) recommend recognition of three genera for the chipmunks: *Tamias* for *T. striatus*, *Eutamias* for *E. sibiricus*, and *Neotamias* for the remaining chipmunks. Our MP and MCMC analyses are essentially in agreement with the hypotheses of Piaggio and Spicer (2000, 2001), supporting a monophyletic ancestry for chipmunks (PP = 99%, BP < 50%), monophyly of the genus *Neotamias* (PP = 100%, BP = 100%), and the uniqueness of *Tamias* and *Eutamias*. The MCMC analysis recovered *E. sibiricus* sister to *T. striatus* (PP = 72%), and this pair sister to the remaining chipmunks (*Neotamias*), whereas the MP analysis resolved *E. sibiricus* basal to all chipmunks and *T. striatus* sister to *Neotamias*.

In both MP and MCMC analyses, the genus *Spermophilus* appears paraphyletic with respect to *Cynomys*, *Marmota*, and possibly *Ammospermophilus*, and some current subgeneric designations (*sensu* Hall, 1981) are not natural groupings. The close relationship among these genera is in agreement with previous analyses of fossil (Black, 1963) and immunological (Hafner, 1984) data, which suggest that they are all descendants of the Miocene *Miospermophilus*. Results from our analyses, however, differ substantially from traditional taxonomy with respect to relationships within this group. Hafner (1984) considered *Ammospermophilus* to have diverged first from the remaining marmotines. Our data do not completely contradict this, but suggest that two species of *Spermophilus* (*S. adocetus* + *S. annulatus*) share an exclusive common ancestor with *Ammospermophilus*. Fossil (Black, 1963) and chromosomal (Nadler et al., 1971) evidence suggest that *Cynomys* is nested within *Spermophilus* and closely allied with the subgenus *Spermophilus*. Our analyses agree that *Cynomys* is derived from within the currently prescribed genus *Spermophilus*, but reject a close relationship with the subgenus *Spermophilus*.

Our phylogenetic hypotheses demonstrate the multiple polyphyletic origins of the genus *Spermophilus*. As currently defined, this genus contains a minimum of seven well-supported, independent clades in the MCMC analysis (Fig. 3, clades S-1 to S-7). Despite some rearrangements among the deepest internodes within the Marmotini, the MP analysis resolved the same seven clades (Fig. 1, clades S-1 to S-7). Clade S-1 (PP = 100%, BP = 99%) includes two members of subgenus *Otospermophilus*, *S. adocetus* and *S. annulatus*, and is sister to the antelope squirrels (PP = 60%, BP = 52%). Clade S-2 (PP = 99%, BP = 90%) is made up of the New World members of the subgenus *Spermophilus* plus the Old World *S. (S.) undulatus*. Clade S-3 (PP = 100%, BP = 100%) includes two members of subgenus *Ictidomys*, *S. mexicanus* and *S. tridecemlineatus*. Clade S-4 includes the only member of subgenus *Poliocitellus*, *S. franklinii*. Clade S-5 (PP = 100%, BP = 87%) includes the

remaining members of subgenus *Ictidomys*, *S. perotensis* and *S. pilosoma*, and both members of subgenus *Xerospermophilus*, *S. mohavensis* and *S. tereticaudus*, and is sister to the prairie dogs (PP = 94%, BP = 57%). Clade S-6 (PP = 100%, BP = 92%) is made up of the Old World members of subgenus *Spermophilus*, except *S. undulatus*. Clade S-7 (PP = 100%, BP = 88%) includes the remaining members of subgenus *Otospermophilus*; *S. atricapillus*, *S. beechyi* and *S. variegatus*; and the three members of subgenus *Callospermophilus*; *S. lateralis*, *S. madrensis* and *S. saturatus*; and is sister to the marmots in the MCMC analysis (PP = 78%, MP does not support).

The taxonomic status of the genus *Spermophilus* is clearly a major problem given our conclusion that it appears to include a total of at least seven independent lineages. All of these clades (S-1 to S-7 in Figs. 1 and 3) are well supported and are sensible candidates for unique generic designations. Given the complex taxonomic history of the Sciuridae, the problem of generic taxonomy requires a comprehensive treatment including a redefinition of available generic names. The type species of the genus *Spermophilus* Cuvier 1825 is *S. citellus*. Thus, the name *Spermophilus* is appropriate for the species included in the Old World clade of the subgenus *Spermophilus* (clade S-6 on Figs. 1 and 3; BP = 92%, PP = 100%). These include *S. citellus*, *S. dauricus*, *S. erythrogegens*, *S. fulvus*, *S. major*, *S. musicus*, *S. pallicauda*, *S. pygmaeus*, *S. relictus*, *S. suslicus*, and *S. xanthoprimum*.

In summary, our results have demonstrated that the current taxonomy of the Sciuridae poorly represents the phylogeny at several levels. Although many taxonomic problems remain, potential solutions are finally becoming apparent. The analyses of Mercer and Roth (2003) and Steppan et al. (in press) provided the basis for a new classification at the subfamilial and tribal levels, but well-supported differences among our phylogenetic hypotheses and theirs must be addressed by future studies that incorporate both a diversity of genetic loci and a more comprehensive representation of taxa. At the tribal level, their analyses found the Funambulini and the Tamiasciurini polyphyletic, and both their analyses and ours show that *Sciurillus* does not belong within the tribe Sciurini. At the genus level, the widely polyphyletic origins of *Spermophilus* suggest a need for further studies that similarly densely sample large genera within this diverse family. Finally, although not the focus of this study, the presence of several species that do not appear monophyletic in either of our reconstructions underscores the need for careful consideration of alpha taxonomy among squirrels.

Acknowledgments

We thank Sue McLaren of the Carnegie Museum of Natural History for providing *Xerus rutilus* tissues, Jane

Waterman of the University of Central Florida for providing *X. inauris* tissues, and Tatsuo Oshida of Tunghai University for providing *Callosciurus* sequences. Support was provided by a grant from the University of Central Florida Department of Biology. Jane Waterman, Tiffany Doan, Robert Ruggiero, and two anonymous reviewers are thanked for helpful suggestions and comments.

References

- Adkins, R.M., Gelke, E.L., Rowe, D., Honeycutt, R.L., 2001. Molecular phylogeny and divergence time estimates for major rodent groups: evidence from multiple genes. *Mol. Biol. Evol.* 18, 777–791.
- Anthony, H.E., Tate, G.H.H., 1935. Notes on South American Mammalia—No. I. *Sciurillus*. *Am. Mus. Novit.* 780, 1–13.
- Arbogast, B.S., Browne, R.A., Weigl, P.D., 2001. Evolutionary genetics and Pleistocene biogeography of North American tree squirrels (*Tamiasciurus*). *J. Mamm.* 82, 302–319.
- Ball, S.S., Roth, V.L., 1995. Jaw muscles of New World squirrels. *J. Morphol.* 224, 265–291.
- Black, C.C., 1963. A review of the North American Tertiary Sciuridae. *Bull. Mus. Comp. Zool.* 130, 109–248.
- Black, C.C., 1972. Holarctic evolution and dispersal of squirrels (Rodentia: Sciuridae). *Evol. Biol.* 6, 305–322.
- Bryant, M.D., 1945. Phylogeny of nearctic sciuridae. *Am. Midl. Nat.* 33, 257–390.
- DeBry, R.W., Sagel, R.M., 2001. Phylogeny of Rodentia (Mammalia) inferred from the nuclear-encoded gene IRBP. *Mol. Phylogenet. Evol.* 19, 290–301.
- Ellerman, J.R., Hayman, R.W., Holt, G.W.C., 1940. The Families and Genera of Living Rodents. Trustees of the British Museum, London, England.
- Ellis, L.S., Maxson, L.R., 1979. Evolution of the chipmunk genera *Eutamias* and *Tamias*. *J. Mamm.* 60, 331–334.
- Ellis, L.S., Maxson, L.R., 1980. Albumin evolution within New World squirrels (Sciuridae). *Am. Midl. Nat.* 104, 57–62.
- Engesser, B., 1979. Relationships of some insectivores and rodents from the Miocene of North America and Europe. *Bull. Carnegie Mus. Nat. Hist.* 14, 1–45.
- Felsenstein, J., 1968. Statistical Inference and the Estimation of Phylogenies. University of Chicago, Chicago.
- Forsyth Major, C.J., 1893. On some Miocene squirrels, with remarks on the dentition and classification of the Sciurinae. *Proc. Zool. Soc. Lond.*, 179–215.
- Griffiths, C.S., 1997. Correlation of functional domains and rates of nucleotide substitution in cytochrome *b*. *Mol. Phylogenet. Evol.* 7, 352–365.
- Hafner, D.J., 1984. Evolutionary relationships of the Nearctic Sciuridae. In: Murie, J., Michener, G. (Eds.), *The Biology of Ground-Dwelling Squirrels: Annual Cycles, Behavioral Ecology, and Sociality*. University of Nebraska Press, Lincoln, pp. 3–23.
- Hafner, M., Barkley, L., Chupasko, J., 1994. Evolutionary genetics of New World tree squirrels (tribe Sciurini). *J. Mamm.* 75, 102–109.
- Hall, E.R., 1981. *The Mammals of North America*. Wiley, New York.
- Hall, E.R., Kelson, K.R., 1959. *The Mammals of North America*. The Ronald Press Company, New York.
- Hight, M.E., Goodman, M., Prychodko, W., 1974. Immunological studies of the Sciuridae. *Syst. Zool.* 23, 12–25.
- Hoffmann, R.S., Anderson, C.G., Thorington Jr., R.W., Heaney, L.R., 1993. Family Sciuridae. In: Wilson, D.E., Reeder, D.M. (Eds.), *Mammal Species of the World: A Taxonomic and Geographic Reference*. Smithsonian Institution Press, Washington, DC, pp. 419–466.
- Howell, N., 1989. Evolutionary conservation of protein regions in the proton-motive cytochrome *b* and their possible roles in redox catalysis. *J. Mol. Evol.* 29, 157–169.
- Huchon, D., Catzeflis, F.M., Douzery, E.J., 1999. Molecular evolution of the nuclear von Willebrand factor gene in mammals and the phylogeny of rodents. *Mol. Biol. Evol.* 16, 577–589.
- Huchon, D., Catzeflis, F.M., Douzery, E.J., 2000. Variance of molecular datings, evolution of rodents and the phylogenetic affinities between Ctenodactylidae and Hystricognathi. *Proc. R. Soc. Lond. B Biol. Sci.* 267, 393–402.
- Huelsenbeck, J.P., Crandall, K.A., 1997. Phylogeny estimation and hypothesis testing using maximum likelihood. *Annu. Rev. Ecol. Syst.* 28, 437–466.
- Huelsenbeck, J.P., Ronquist, F., 2001. MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* 17, 754–755.
- Huelsenbeck, J.P., Larget, B., Miller, R.E., Ronquist, F., 2002. Potential applications and pitfalls of bayesian inference of phylogeny. *Syst. Biol.* 51, 672–688.
- Irwin, D., Kocher, T., Wilson, A., 1991. Evolution of the cytochrome *b* gene in mammals. *J. Mol. Evol.* 32, 37–55.
- James, G.T., 1963. Paleontology and nonmarine stratigraphy of the Cuyama Valley Badlands, California. Part I. Geology, faunal interpretations, and systematic descriptions of Chiroptera, Insectivora, and Rodentia. *Univ. Calif. Publ. Geol. Sci.* 45, 1–171.
- Jameson Jr., E.W., 1999. Host–ectoparasite relationships among North American chipmunks. *Acta Theriol.* 44, 225–231.
- Kimura, M., 1986. The role of compensatory neutral mutations in molecular evolution. *J. Genet.* 64, 7–19.
- Krajewski, C., King, D.G., 1996. Molecular divergence and phylogeny: rates and patterns of cytochrome *b* evolution in cranes. *Mol. Biol. Evol.* 13, 21–30.
- Lavocat, R., Parent, J.-P., 1985. Phylogenetic analysis of middle-ear features in fossil and living rodents. In: Lockett, W.P., Hartenberger, J.-L. (Eds.), *Evolutionary Relationships Among Rodents: A Multidisciplinary Analysis*. Plenum Press, New York, pp. 333–354.
- Layne, J.N., 1952. The *os genitale* of the red squirrel, *Tamiasciurus*. *J. Mamm.* 33, 457–459.
- Levenson, H., Hoffman, R.S., Nadler, C.F., Deutsch, L., Freeman, S.D., 1985. Systematics of the Holarctic chipmunks (*Tamias*). *J. Mamm.* 66, 219–242.
- McKenna, M.C., Bell, S.K., 1997. *Classification of Mammals Above the Species Level*. Columbia University Press, New York.
- Mein, P., 1970. Les sciuropteres (Mammalia, Rodentia) Neogenes d'Europe occidentale. *Geobios* 3, 7–77.
- Mercer, J.M., Roth, V.L., 2003. The effects of Cenozoic global change on squirrel phylogeny. *Science* 299, 1568–1572.
- Montgelard, C., Bentz, S., Tirard, C., Verneau, O., Catzeflis, F.M., 2002. Molecular systematics of Sciurognathi (Rodentia): the mitochondrial cytochrome *b* and 12S rRNA genes support the Anomaluroidea (Pedetidae and Anomaluridae). *Mol. Phylogenet. Evol.* 22, 220–233.
- Moore, J.C., 1959. Relationships among the living squirrels of the Sciurinae. *Bull. Am. Mus. Nat. Hist.* 118, 159–206.
- Moore, J.C., 1961. Geographic variation in some reproductive characteristics of diurnal squirrels. *Bull. Am. Mus. Nat. Hist.* 122, 1–32.
- Mossman, H.W., Lawlah, J.W., Bradley, J.A., 1932. The male reproductive tract of the Sciuridae. *Am. J. Anat.* 51, 89–155.
- Nadler, C.F., 1964. Contributions of chromosomal analysis to the systematics of North American chipmunks. *Am. Midl. Nat.* 72, 298–312.
- Nadler, C.F., Hoffmann, R.S., Lay, D.M., 1969. Chromosomes of the Asian chipmunk *Eutamias sibiricus* Laxmann (Rodentia: Sciuridae). *Experientia* 25, 868–869.
- Nadler, C.F., Hoffmann, R.S., Pizzimenti, J.J., 1971. Chromosomes and serum proteins of prairie dogs and a model of *Cynomys* evolution. *J. Mamm.* 52, 545–555.

- Nadler, C.F., Hoffmann, R.S., Honacki, J.H., Pozin, D., 1977. Chromosomal evolution in chipmunks, with special emphasis on A and B karyotypes of the subgenus *Neotamias*. *Am. Midl. Nat.* 98, 343–353.
- Nedbal, M.A., Honeycutt, R.L., Schlitter, D.A., 1996. Higher-level systematics of rodents (Mammalia: Rodentia): Evidence from the mitochondrial 12S rRNA gene. *J. Mamm. Evol.* 3, 201–237.
- Nicholas, K.B., Nicholas Jr., H.B., 1997. Genedoc: a tool for editing and annotating multiple sequence alignments. Distributed by the authors.
- Nielsen, R., 1997. Site-by-site estimation of the rate of substitution and the correlation of rates in mitochondrial DNA. *Syst. Biol.* 46, 346–353.
- Oshida, T., Masuda, R., 2000. Phylogeny and zoogeography of six squirrel species of the genus *Sciurus* (Mammalia, Rodentia), inferred from cytochrome *b* gene sequences. *Zool. Sci.* 17, 405–410.
- Oshida, T., Masuda, R., Yoshida, M., 1996. Phylogenetic relationships among Japanese species of the family Sciuridae (Mammalia, Rodentia), inferred from nucleotide sequences of mitochondrial 12S ribosomal RNA genes. *Zool. Sci.* 13, 615–620.
- Oshida, T., Lin, L.K., Masuda, R., Yoshida, M.C., 2000. Phylogenetic relationships among Asian species of *Petaurista* (Rodentia, Sciuridae), inferred from mitochondrial cytochrome *b* gene sequences. *Zool. Sci.* 17, 123–128.
- Oshida, T., Yasuda, M., Endo, H., Hussein, N.A., Masuda, R., 2001. Molecular phylogeny of five squirrel species of the genus *Callosciurus* (Mammalia, Rodentia) inferred from cytochrome *b* gene sequences. *Mammalia* 65, 473–482.
- Piaggio, A.J., Spicer, G.S., 2000. Molecular phylogeny of the chipmunk genus *Tamias* based on the mitochondrial cytochrome oxidase subunit II gene. *J. Mamm. Evol.* 7, 147–166.
- Piaggio, A.J., Spicer, G.S., 2001. Molecular phylogeny of the chipmunks inferred from mitochondrial cytochrome *b* and cytochrome oxidase II gene sequences. *Mol. Phylogenet. Evol.* 20, 335–350.
- Pocock, R.I., 1923. The classification of the Sciuridae. *Proc. Zool. Soc. Lond.* 1923, 209–246.
- Posada, D., Crandall, K., 1998. MODELTEST: testing the model of DNA substitution. *Bioinformatics* 14, 817–818.
- Posada, D., Crandall, K.A., 2001. Selecting the best-fit model of nucleotide substitution. *Syst. Biol.* 50, 580–601.
- Saiki, R., Gelfand, D.H., Stoffl, S., Scharf, S., Higuchi, T., Horn, G., Mullis, K.B., Erlich, H., 1988. Primer-directed enzymatic amplification of DNA with thermostable DNA polymerase. *Science* 239, 487–491.
- Sambrook, J., Russell, D.W., 2001. *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Schöniger, M., Von Haeseler, A., 1994. A stochastic model for the evolution of autocorrelated DNA sequences. *Mol. Phylogenet. Evol.* 3, 240–247.
- Simpson, G.G., 1945. Principles of classification and a classification of mammals. *Bull. Am. Mus. Nat. Hist.* 85, 1–350.
- Smith, A.B., 1994. Rooting molecular trees: problems and strategies. *Biol. J. Linnean Soc.* 51, 279–292.
- Steppan, S.J., Akhverdyan, M.R., Lyapunove, E.A., Fraser, D.G., Vorontsov, N.N., Hoffmann, R.S., Braun, M.J., 1999. Molecular phylogeny of the marmots (Rodentia: Sciuridae): tests of evolutionary and biogeographic hypotheses. *Syst. Biol.* 48, 715–734.
- Steppan, S.J., Storz, B.L., Hoffmann, R.S., in press. Nuclear DNA phylogeny of the squirrels (Mammalia: Rodentia) and the evolution of arboreality from *c-myc* and *RAG1*. *Mol. Phylogenet. Evol.*
- Swofford, D.L., 2002. *PAUP*. Phylogenetic Analysis Using Parsimony (* and Other Methods)*. Sinauer Associates, Sunderland, MA.
- Tavaré, S., 1986. Some probabilistic and statistical problems in the analysis of DNA sequences. In: Miura, R.M. (Ed.), *Some Mathematical Questions in Biology/DNA Sequence Analysis*. American Mathematical Society, Providence, RI, pp. 57–86.
- Thorington Jr., R.W., 1984. Flying squirrels are monophyletic. *Science* 225, 1048–1050.
- Thorington Jr., R.W., Darrow, K., 1996. Jaw muscles of Old World squirrels. *J. Morphol.* 230, 145–165.
- Wade, O., Gilbert, P.T., 1940. The baculum of some Sciuridae and its significance in determining relationships. *J. Mamm.* 21, 52–63.
- Wahlert, J.H., 1985. Cranial foramina of rodents. In: Lockett, W.P., Hartenberger, J.-L. (Eds.), *Evolutionary Relationships Among Rodents: A Multidisciplinary Analysis*. Plenum Press, New York, pp. 311–332.
- Wettstein, P.J., Strausbauch, M., Lamb, T., States, J., Chakraborty, R., Jin, L., Riblet, R., 1995. Phylogeny of six *Sciurus aberti* subspecies based on nucleotide sequences of cytochrome *b*. *Mol. Phylogenet. Evol.* 4, 150–162.
- White, J.W., 1953. Genera and subgenera of chipmunks. *Univ. Kansas Publ. Mus. Nat. Hist.* 5, 543–561.
- Yang, Z., 1993. Maximum-likelihood estimation of phylogeny from DNA sequences when substitution rates differ over sites. *Mol. Biol. Evol.* 10, 1396–1401.
- Yang, Z., 1995. A space-time process model for the evolution of DNA sequences. *Genetics* 139, 993–1005.